



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# Large-eddy simulation of very large kinetic and magnetic Reynolds number isotropic magnetohydrodynamic turbulence using a spectral subgrid model

T. Gomez, P. Sagaut, O. Schilling, Y. Zhou

July 7, 2006

Physics of Fluids

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

# Large-eddy simulation of very large kinetic and magnetic Reynolds number isotropic magnetohydrodynamic turbulence using a spectral subgrid model

Th. Gomez<sup>(a,)</sup>, P. Sagaut<sup>(a)</sup>, O. Schilling<sup>(b)</sup> & Y. Zhou<sup>(b)</sup>

(a) Laboratoire de Modélisation en Mécanique,  
Université Pierre et Marie Curie - Paris 6,  
4 place Jussieu - case 162,

F-75252 Paris cedex 5, France

(b) University of California, Lawrence Livermore National Laboratory,  
Livermore, California 94551, USA

PACS numbers: 47.27.ep, 52.30.cv

A spectral subgrid-scale eddy viscosity and magnetic resistivity model based on the eddy-damped quasi-normal Markovian (EDQNM) spectral kinetic and magnetic energy transfer presented in [12] is used in large-eddy simulation (LES) of large kinetic and magnetic Reynolds number magnetohydrodynamic (MHD) turbulence. The proposed model is assessed via *a posteriori* tests on three-dimensional, incompressible, isotropic, non-helical, freely-decaying MHD turbulence at asymptotically large Reynolds numbers. Using LES with an initial condition characterized by an Alfvén ratio of kinetic to magnetic energy  $r_A$  equal to unity, it is shown that the kinetic energy spectrum  $E_K(k)$  and magnetic energy spectrum  $E_M(k)$  exhibit Kolmogorov  $-5/3$  inertial subrange scalings in the LES, consistent with the EDQNM model.

PACS numbers:

Numerical simulations for investigating the physics of magnetohydrodynamic (MHD) turbulence are of the greatest interest. As most astrophysical and geophysical plasmas (e.g. the liquid core of the Earth, accretion disks, and star-forming molecular clouds) cannot be directly investigated experimentally, numerical simulations are the method of choice for studying the properties of such plasmas. In the field of applied and fundamental research related to engineering and physical sciences, most flows are highly turbulent. Even with contemporary supercomputing capability, direct numerical simulation (DNS) remains limited to turbulent flows with modest Reynolds numbers. Large-eddy simulation (LES) overcomes this computational limitation by computing only the largest resolved scales and modeling the effects of the subgrid scales on these resolved scales using physical arguments to approximate turbulent energy dissipation and backscatter [7].

LES is based on the separation of the spatial scales of motion: the resolved scales are directly computed by solving partial differential equations consisting of the usual governing equations supplemented with a subgrid-scale term which account for the effects of the unresolved scales of motion. Two different types of subgrid-scale models can be used for isotropic turbulence: physical space models for finite-element, finite-volume and finite-difference methods, or spectral models for (pseudo)-spectral methods. Physical space subgrid-scale models for MHD flows are typically extensions of the models introduced in the pure hydrodynamic case, but are used

along with non-spectral numerical methods which are less accurate for the resolution of the small scales than spectral methods. The numerical error induced by these schemes is often very large in the vicinity of the cutoff wave number, resulting in a poor estimate of the energy transfer across the cutoff scale. On the other hand, spectral methods are more accurate at the small scales where the interscale dynamics can be well captured, yielding an accurate prediction of the subgrid-scale energy transfer. This remark is well founded, as shown by the energy transfer cusp, observed near the cutoff in direct numerical simulations. This cusp is underestimated in physical space eddy viscosity-type subgrid models with model constants computed dynamically, as shown in [4] by comparison with filtered DNS data, whereas it is clearly present in the spectral subgrid model as shown, for example, in the small Prandtl number MHD subgrid-scale model of Ponty *et al.* [6].

Historically, subgrid-scale models for MHD turbulence expressed in physical space have been constructed by extending the usual non-magnetic models to the case of electrically-conducting fluids [11]. MHD gradient-diffusion type subgrid-scale models have been assessed for isotropic turbulence at magnetic Prandtl number  $Pr_m = 1$  [1, 4] and at small magnetic Prandtl number [3]. In the anisotropic case, this type of subgrid-scale model was used for turbulent channel flow at small magnetic Prandtl number and kinetic Reynolds number  $Re = 29000$  [10]. Additionally, a physical space subgrid-scale model based on the ideal cross-helicity invariant of

MHD turbulence has been developed [4, 5] to account for the cross-helicity inverse cascade in the energy transfer. These models appear to capture the principal features of incompressible homogeneous MHD turbulence at a computational cost orders of magnitude lower than a DNS.

More recently, a spectral subgrid-scale model was developed for MHD turbulence [12] using an analysis of the eddy-damped quasi-normal Markovian (EDQNM) subgrid-scale kinetic and magnetic energy transfers in isotropic turbulence; however, this model has not yet been assessed using LES. This model was derived by applying a sharp Fourier cutoff filter to the spectral energy transfer equations, generalizing the analysis in [8] to MHD turbulence. In the case of small magnetic Prandtl number statistically-stationary turbulence with an external constant large-scale magnetic field, Ponty *et al.* [6] empirically constructed a Chollet-Lesieur [2] type spectral subgrid model. In these small magnetic Prandtl number simulations, the magnetic field fluctuations are fully resolved and the subgrid velocity fluctuations are modeled using LES. The results were found to be in good agreement with existing experimental data for large kinetic Reynolds number.

In nondimensional form, the unforced incompressible MHD equations (with unit constant density) are written as

$$\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} = (\nabla \times \mathbf{b}) \times \mathbf{b} - \nabla p + \nu_T \nabla^2 \mathbf{v} \quad (1)$$

$$\begin{aligned} \frac{\partial \mathbf{b}}{\partial t} &= \nabla \times (\mathbf{u} \times \mathbf{b}) + \eta_T \nabla^2 \mathbf{b} \\ \nabla \cdot \mathbf{v} &= 0, \quad \nabla \cdot \mathbf{b} = 0, \end{aligned} \quad (2)$$

where  $\mathbf{v}$  is the fluid velocity,  $\mathbf{b}$  is the magnetic field,  $p$  is the hydrodynamic pressure, and  $\nu_T$  and  $\eta_T$  are the eddy viscosity and magnetic resistivity, respectively. Only infinite kinetic and magnetic Reynolds numbers are considered, so that the only dissipative mechanism in the system is the EDQNM-based spectral eddy viscosity and magnetic resistivity.

As shown in [12], the eddy viscosity and resistivity are much larger than the backscatter counterpart, except very close to the cusp  $k/k_c \lesssim 1$  where the ratio is still larger than two. Therefore, the effect of backscatter on the total amount of subgrid transfer will be neglected in the present study. To parameterize the eddy viscosity  $\nu_T$  and magnetic resistivity  $\eta_T$  for use in the LES, these functions are fit as a function of  $k/k_c$  from the numerical values obtained from the EDQNM closure computations [9, 12]. The form used for  $\nu^+(k|k_c; t)$  and  $\eta^+(k|k_c; t)$  is

$$\begin{aligned} \nu^+(k|k_c; t) &= \\ &\left[ 0.0023398 + 0.0392187 \left( \frac{k}{k_c} \right) + 3.73676 \left( \frac{k}{k_c} \right)^4 \right] \\ &\times \exp \left[ 4.67997 - 13.121 \left( \frac{k}{k_c} \right) + 6.97447 \left( \frac{k}{k_c} \right)^2 \right] \\ \eta^+(k|k_c; t) &= \end{aligned}$$

$$\begin{aligned} &\left[ 0.00267451 + 0.0509853 \left( \frac{k}{k_c} \right) + 3.61643 \left( \frac{k}{k_c} \right)^4 \right] \\ &\times \exp \left[ 4.51932 - 13.4592 \left( \frac{k}{k_c} \right) + 7.37105 \left( \frac{k}{k_c} \right)^2 \right] \end{aligned}$$

where  $k_c$  is the cutoff wavenumber, and the eddy viscosity and magnetic resistivity are conventionally normalized as

$$\nu_T(k|k_c; t) = \nu^+(k|k_c; t) \sqrt{\frac{E_K(k_c, t)}{k_c}} \quad (3)$$

$$\eta_T(k|k_c; t) = \eta^+(k|k_c; t) \sqrt{\frac{E_M(k_c, t)}{k_c}}. \quad (4)$$

The kinetic and magnetic energy are

$$E_K(t) = \frac{1}{2} \int_V \mathbf{v}^2 d^3x, \quad (5)$$

$$E_M(t) = \frac{1}{2} \int_V \mathbf{b}^2 d^3x, \quad (6)$$

respectively, with  $V$  the volume of the domain. The kinetic to magnetic energy (Alfvén) ratio  $r_A = E_K(t)/E_M(t)$  characterizes the energy ratio at the large scales of the flow. The kinetic and magnetic enstrophies are

$$\Omega_K(t) = \frac{1}{2} \int_V (\nabla \times \mathbf{v})^2 d^3x, \quad (7)$$

$$\Omega_M(t) = \frac{1}{2} \int_V (\nabla \times \mathbf{b})^2 d^3x, \quad (8)$$

respectively. The kinetic to magnetic enstrophy ratio  $r_B = \Omega_K(t)/\Omega_M(t)$  characterizes the energy ratio at the small scales of the flow.

A three-dimensional freely-decaying LES is performed in a  $(2\pi)^3$  periodic computational domain using uniform  $N^3 = 128^3$  and  $N^3 = 64^3$  computational grids. The dissipative terms in the kinetic and magnetic equations are given respectively by (3) and (4).

The initial fields are constructed as a sum of large-scale Fourier modes with random amplitudes and phases. The kinetic and magnetic energy are both normalized to 0.5 ( $r_A = 1$ ) and the initial cross-correlation  $E_c = 2\langle |\mathbf{u} \cdot \mathbf{b}| \rangle / \langle \mathbf{u}^2 + \mathbf{b}^2 \rangle$  is approximately 0.2.

The simulation is performed using a pseudospectral code to solve the incompressible three-dimensional equations (1)–(2) with desaliasing performed according to the 2/3 rule. A third-order Runge-Kutta scheme is used for the time integration with a constant timestep  $\Delta t = 2.5 \times 10^{-3}$ .

The temporal evolution of the kinetic and magnetic energy exhibits a self-similar decay between  $t = 5$  and 18 as shown in Fig. 2. During this period, the kinetic energy decreases slightly more rapidly according to  $E_K(t) \sim t^{-1.4}$  than the magnetic energy  $E_M(t) \sim t^{-1.3}$ . Figures (5) and (6) show the kinetic and magnetic spectra




FIG. 1: The eddy viscosity (top) and magnetic resistivity (bottom): fit (—) and values from EDQNM closure calculation (+).




FIG. 2: Temporal evolution of the kinetic (—) and magnetic (---) energy (top) and of the enstrophy (bottom) between  $t = 0$  and 18.

at times  $t = 7.5, 8.5, 9.5, 10.5, 11.5$  and  $12.5$ . An inertial subrange with slope  $-5/3$  is observed for both spectra, consistent with the EDQNM model [12]. The kinetic and magnetic spectra are shown at different times to show that the simulations are well converged. Between  $t = 0$  and 18, the large-scale ratio  $r_A$  decreases from 1 to 0.4, whereas the small-scale ratio  $r_B$  is stabilized near 0.7 between  $t = 2.5$  and 18. This ratio variation in the freely-decaying turbulent case shows that the EDQNM-based LES method could possibly be improved by considering eddy viscosities and magnetic resistivities depending on the Alfvén ratio even if the small-scale ratio  $r_B$  remains approximately constant during the course of the simulation.

The spectral subgrid-scale eddy viscosity and resistivity obtained from EDQNM closure calculations [12] have been parameterized and used to perform LES of three-dimensional, isotropic, non-helical, incompressible, magnetohydrodynamic turbulent flow for quasi-equipartition of energy, *i.e.*  $r_A \sim 1$ . This subgrid-scale model is robust, as the LES predict  $-5/3$  inertial subrange kinetic




FIG. 3: Temporal evolution of the kinetic to magnetic energy ratio  $r_A$  (—) and of the kinetic to magnetic enstrophy ratio  $r_B$  (---) between  $t = 0$  and 18.




FIG. 4: Temporal evolution of the correlation  $E_c = 2\langle |\mathbf{u} \cdot \mathbf{b}| \rangle / \langle \mathbf{u}^2 + \mathbf{b}^2 \rangle$  between  $t = 0$  and 18.




FIG. 5: Log-log plot of the kinetic energy spectrum  $E_K(k, t)$  normalized by  $E_K(t)$ :  $N^3 = 128^3$  (—) and  $N^3 = 64$  (+).

and magnetic energy spectra, even when the Alfvén ratio decreases from 1 to 0.4 as shown in the simulations.

#### Acknowledgements

This work was performed under the auspices of the U.S. Department of Energy by the University of California Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

- 
- [1] O. Agullo, W. Müller, B. Knaepen, and D. Carati. Large eddy simulation of decaying magnetohydrodynamic turbulence with dynamic subgrid-modeling. *Phys. of Plasma*, 8(7):3502–3505, 2001.
  - [2] J.-P. Chollet and M. Lesieur. Parametrization of small scales of three-dimensional isotropic turbulence utilizing spectral closures. *J. Atmos. Sci.*, 38:2747–2757, 1981.
  - [3] B. Knaepen and P. Moin. large-eddy simulation of conductive flows at low magnetic reynolds number. *Phys. of Fluids*, 16(5):1255–1261, 2004.
  - [4] W. Müller and D. Carati. Dynamic gradient-diffusion subgrid models for incompressible magnetohydrodynamic turbulence. *Phys. of Plasma*, 9(3):824–834, 2002.
  - [5] W. Müller and D. Carati. Large-eddy simulation for magnetohydrodynamics turbulence. *Comp. Phys. Com.*, 147:544–547, 2002.
  - [6] Y. Ponty, H. Politano, and J.F. Pinton. Simulation of induction at low magnetic prandtl number. *Phys. Rev. Lett.*, 92(14):144503–1–144503–4, 2004.
  - [7] P. Sagaut. *Large eddy simulation for incompressible flows*. Scientific computation series. Springer-Verlag, 2005.
  - [8] Oleg Schilling and Ye Zhou. Analysis of spectral eddy viscosity and backscatter in incompressible, isotropic turbulence using statistical closure theory. *Phys. of Fluids*, 14(3):1244–1258, 2002.
  - [9] Oleg Schilling and Ye Zhou. Triadic energy transfers in non-helical magnetohydrodynamic turbulence. *J. Plasma Physics*, 68:389–406, 2002.
  - [10] Y. Shimomura. Large eddy simulation of magnetohydrodynamic turbulent channel flows under a uniform magnetic field. *Phys. Fluids A*, 3(12):3098–3106, 1991.
  - [11] Y. Zhou and G. Vahala. Aspects of subgrid modeling and large-eddy simulation of magnetohydrodynamic tur-




FIG. 6: Log-log plot of the magnetic energy spectrum  $E_M(k, t)$  normalized by  $E_M(t)$ :  $N^3 = 128^3$  (—) and  $N^3 = 64$  (+).

- bulence. *J. Plasma Physics*, 45:239–249, 1991.
- [12] Ye Zhou, Oleg Schilling, and S. Ghosh. Subgrid scale and backscatter model for magnetohydrodynamic turbulence based on closure theory: theoretical formulation. *Phys. Rev. E*, 66:026309–1–026309–6, 2002.